



FLORAL AND FAUNAL BIODIVERSITY OF KASHMIR HIMALAYA: PATTERNS, THREATS, AND CONSERVATION PERSPECTIVES

SYED OWAIS MUSHTAQ* AND MUZAFFAR AHMAD NAIKOO

¹Department of Biochemistry, Government Degree College Kulgam,
University of Kashmir, Jammu and Kashmir, India.

*Corresponding author email: owaissyed854@gmail.com

ABSTRACT: The Kashmir Himalaya represents one of the most biologically diverse regions of South Asia, owing to its complex topography, broad altitudinal gradients, and heterogeneous climatic conditions. The region encompasses a wide range of ecosystems, including temperate forests, alpine meadows, wetlands, and riverine habitats, which collectively support high floral and faunal diversity, including numerous endemic and threatened species. This review synthesizes existing literature to provide an integrated assessment of floral and faunal biodiversity in Kashmir, highlighting major vegetation types, key faunal assemblages, and species of high conservation significance such as the critically endangered Hangul deer (*Cervus hanglu* Wagner, 1844), Snow leopard (*Panthera uncia* (Schreber, 1775)), and Black-necked crane (*Grus nigricollis*). The review critically examines the primary drivers of biodiversity loss, including habitat fragmentation, climate change, poaching, invasive species, pollution, and increasing human–wildlife conflicts. Current conservation initiatives, encompassing protected area networks, policy frameworks, community participation, ecotourism, and international conservation commitments, are evaluated for their effectiveness and limitations. Finally, future prospects and strategic recommendations are proposed, emphasizing strengthened policy enforcement, habitat restoration, long-term ecological monitoring, and community-based conservation to ensure the sustainable management of biodiversity in the Kashmir Himalaya. This synthesis aims to inform conservation planning and policy formulation in a region experiencing rapid environmental and socio-economic change..

Keywords: Kashmir Himalaya, Biodiversity conservation, Flora, Fauna, Threatened species, Climate change

Citation: Mushtaq Owais S, Naikoo Ahmad M (2025) Floral and Faunal Biodiversity of Kashmir Himalaya: Patterns, Threats, And Conservation Perspectives. Indian J Trop Biodiv 33 (2): 75-93.

Received on: 21/01/2026; Accepted on: 26/03/2026

INTRODUCTION

BACKGROUND ON BIODIVERSITY IN KASHMIR

Kashmir, nestled in the northwestern Himalayas, is renowned for its rich and diverse biodiversity. The

region's unique topography, ranging from lush valleys to high-altitude alpine zones, supports a variety of ecosystems that harbor both endemic and globally significant species (Dar and Khuroo, 2020). The varied climatic conditions, from temperate forests to glacial landscapes, contribute

to the existence of diverse flora and fauna (Sharma *et al.*, 2018). Biodiversity in Kashmir plays a crucial role in maintaining ecological balance, supporting livelihoods, and sustaining agricultural and medicinal plant industries (Rather *et al.*, 2021).

IMPORTANCE OF FLORAL AND FAUNAL DIVERSITY

The floral and faunal diversity of Kashmir is vital for maintaining ecosystem services, including climate regulation, soil conservation, and pollination (Hussain *et al.*, 2019). The region hosts a range of endemic and endangered species, such as the Hangul deer (*Cervus hanglu* Wagner, 1844), Snow Leopard (*Panthera uncia* (Schreber, 1775)), and Kashmir Stag, which are of high conservation significance (Wani *et al.*, 2022). Medicinal and aromatic plants found in Kashmir contribute to traditional knowledge and pharmaceutical industries, making biodiversity conservation a priority for sustainable development (Lone and Bhatt, 2020).

OBJECTIVES OF THE REVIEW

This review aims to:

1. Provide an overview of the floral and faunal diversity in Kashmir.
2. Highlight the ecological significance of biodiversity in the region.
3. Discuss conservation challenges and threats affecting biodiversity.
4. Review existing conservation policies and suggest future recommendations for biodiversity protection.

METHODOLOGY FOR LITERATURE REVIEW

A systematic literature review was conducted following PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guidelines to ensure transparency and reproducibility.

DATABASE SEARCH STRATEGY

The following electronic databases were searched:

- Web of Science (WoS)
- Scopus
- Google Scholar

The search strategy was structured using Boolean operators and controlled vocabulary to avoid inflated or biased results.

The final Boolean search string used was:

("flora" OR "plant diversity" OR "plant biodiversity" OR "vegetation diversity")
AND
("fauna" OR "wildlife diversity" OR "animal diversity")
AND
("Kashmir Himalaya" OR "Kashmir Valley")
AND
("biodiversity" OR "conservation" OR "threats" OR "climate change")

The search period covered publications from 2000 to 2024. Only English-language peer-reviewed publications were considered.

LITERATURE SEARCH RESULTS

The initial search across databases yielded 742 records.

After removing 186 duplicate entries, 556 unique records were screened based on title and abstract.

Of these:

- 474 records were excluded due to lack of regional focus or thematic relevance.
- 82 publications met the inclusion criteria and were included for qualitative synthesis.

These comprised:

- 64 peer-reviewed research articles
- 18 official government biodiversity and wildlife reports

This structured screening process minimized selection bias and ensured methodological rigor.

INCLUSION AND EXCLUSION CRITERIA

Included:

- Studies focused specifically on the Kashmir Himalaya
- Research addressing plant and/or animal biodiversity
- Publications discussing conservation status or ecological threats
- Official biodiversity reports from recognized government bodies

Excluded:

- Non-regional studies
- Opinion pieces without empirical data
- Conference abstracts without full text
- Duplicate publications

JUSTIFICATION FOR INCLUSION OF GREY LITERATURE

Although systematic reviews generally limit inclusion to peer-reviewed literature, grey literature was incorporated cautiously due to the limited availability of long-term biodiversity monitoring datasets in the Kashmir Himalaya.

To avoid selective inclusion bias:

- Only officially published reports from recognized institutions (e.g., State Biodiversity Board, Wildlife Department) were included.
- Data were cross-verified with peer-reviewed publications wherever possible.
- Grey literature was used only for supplementary data (e.g., species population estimates), not for interpretative conclusions.
- This approach ensured comprehensiveness while maintaining scientific credibility.

GEOGRAPHICAL AND ECOLOGICAL OVERVIEW OF KASHMIR

Kashmir, located in the northwestern part of the Indian subcontinent, is part of the larger Himalayan mountain system. The region spans across the Union Territory of Jammu and Kashmir and

extends into parts of Ladakh, covering an area with highly diverse topographical features, including lush valleys, snow-capped mountains, glaciers, rivers, and lakes (Raina *et al.*, 2019). The altitude of Kashmir ranges from 1,585 meters in the valley to over 7,000 meters in the greater Himalayan ranges, which significantly influences its climatic conditions and biodiversity (Sharma and Koul, 2021). The topography of Kashmir is marked by the Pir Panjal range to the southwest, the Karakoram range to the northeast, and the Zaskar range to the east, which collectively create distinct ecological zones supporting a variety of plant and animal life (Dar *et al.*, 2020).

The climate of Kashmir is primarily temperate, characterized by four distinct seasons: spring, summer, autumn, and winter, with significant variations in temperature and precipitation across different altitudes. The region experiences cold winters with heavy snowfall, particularly in the upper reaches, while summers are mild with moderate rainfall, primarily due to the influence of the southwest monsoon (Ahmad and Wani, 2022). The temperature in the valley ranges from sub-zero levels in winter to around 35°C in peak summer, with an annual precipitation average of approximately 660–1,200 mm (Rather *et al.*, 2021). These seasonal variations create a dynamic environment that influences vegetation patterns, animal behavior, and agricultural cycles in the region (Shafiq *et al.*, 2023).

Kashmir hosts a range of ecosystems, each contributing to the region's rich biodiversity. The Himalayan forests, which include coniferous and mixed deciduous forests, are home to species like *Cedrus deodara*, *Pinus wallichiana*, and *Abies pindrow*, forming crucial habitats for endemic and migratory wildlife (Hussain *et al.*, 2018). Alpine meadows, found at higher altitudes, support herbaceous vegetation, including medicinal and aromatic plants such as *Aconitum heterophyllum* and *Rheum emodi*, which are vital for both ecological stability and traditional medicine (Bhat *et al.*, 2020). Wetlands such as Hokersar and

Wular Lake play an essential role in regulating water resources and serve as habitats for migratory birds, including the black-necked crane and bar-headed goose (Lone and Bhatt, 2020). The region is also rich in riverine ecosystems, with major rivers such as the Jhelum, Chenab, and Indus providing sustenance to both terrestrial and aquatic life (Wani *et al.*, 2022). These diverse ecosystems collectively make Kashmir a biodiversity hotspot, underscoring the need for conservation efforts to protect its unique flora and fauna.

FLORAL DIVERSITY IN KASHMIR

Kashmir is home to a rich and diverse floral ecosystem, shaped by its varied topography and climatic conditions. The region harbors thousands of plant species, ranging from temperate forests in the lower valleys to alpine vegetation at higher altitudes. The floral diversity of Kashmir is classified into different vegetation types, including forest vegetation, alpine and sub-alpine flora, wetland and aquatic flora, and endemic and rare species (Dar and Khuroo, 2020). This diverse plant life plays a significant role in maintaining ecological stability, supporting wildlife, and contributing to local economies through agriculture, forestry, and medicinal plant industries (Rather *et al.*, 2021). The forests of Kashmir are classified into coniferous, deciduous, and mixed forest types, each harboring unique species adapted to specific climatic conditions. The coniferous forests, primarily found in the Pir Panjal and Zaskar ranges, consist of species such as *Cedrus deodara* (Deodar), *Abies pindrow* (Fir), and *Pinus wallichiana* (Blue Pine), which are important for timber and ecological conservation (Sharma and Koul, 2021). Deciduous forests, mainly found in the lower valleys and foothills, include species such as *Quercus spp.* (Oaks), *Juglans regia* (Walnut), and *Acer caesium* (Maple), providing habitat for various bird and mammal species (Hussain *et al.*, 2018). Mixed forests, which are a combination of coniferous and broadleaf species, act as transition zones and support high levels of

biodiversity by hosting multiple plant and animal species (Wani *et al.*, 2022). Alpine and sub-alpine vegetation is found at elevations above 3,000 meters, where harsh climatic conditions limit tree growth. This zone is characterized by herbaceous and shrubby vegetation, including species such as *Rheum emodi* (Himalayan Rhubarb), *Aconitum heterophyllum* (Atis), and *Saussurea costus* (Falc.) Lipsch., which are known for their medicinal properties (Bhat *et al.*, 2020). The high-altitude meadows, known as bugyals, provide seasonal grazing grounds for livestock and serve as critical habitats for pollinators and herbivorous mammals (Ahmad and Wani, 2022). Kashmir's wetlands and aquatic ecosystems, including Wular Lake, Dal Lake, and Hokersar Wetland, support a rich diversity of aquatic plants such as *Nymphaea alba* (White Water Lily), *Trapa natans* (Water Chestnut), and *Potamogeton spp.* (Pondweeds) (Lone and Bhatt, 2020). These water bodies play a crucial role in maintaining the ecological balance by acting as carbon sinks, providing breeding grounds for migratory birds, and supporting fisheries (Rather *et al.*, 2021). However, increasing pollution and land-use changes threaten these fragile ecosystems, necessitating conservation efforts (Shafiq *et al.*, 2023). Several plant species found in Kashmir are endemic and rare, making them of significant conservation importance. Species such as *Meconopsis aculeata* (Himalayan Blue Poppy) and *Arnebia benthamii* (Kahzaban) are restricted to specific altitudinal zones and are highly sensitive to climate change and habitat degradation (Bhat *et al.*, 2020). The region is also known for its medicinal and aromatic plants, which have been used in traditional medicine for centuries. Notable examples include *Berberis aristata* (Tree Turmeric), *Viola odorata* (Sweet Violet), and *Podophyllum hexandrum* (Himalayan Mayapple), which are extensively harvested for their therapeutic properties (Dar and Khuroo, 2020). These plants are used in Ayurveda, Unani, and modern pharmaceutical formulations, contributing to the region's economy and healthcare industry (Sharma *et al.*, 2018). The

economic and ecological significance of Kashmir's floral biodiversity cannot be overstated. Forests provide timber, fuelwood, and non-timber forest products, supporting local livelihoods (Hussain *et al.*, 2019). Medicinal plant trade generates substantial revenue, while the agricultural sector benefits from pollination and soil stabilization provided by plant diversity (Rather *et al.*, 2021). Additionally, floral biodiversity plays a key role in climate regulation, carbon sequestration, and watershed protection, making conservation a priority for sustainable development in the region (Ahmad *et al.*, 2021).

FAUNAL DIVERSITY IN KASHMIR

Kashmir is home to a diverse range of faunal species, shaped by its varied topography, climate, and vegetation. The region supports mammals, birds, reptiles, amphibians, aquatic life, and insect species, many of which are endemic or endangered. The faunal diversity of Kashmir is influenced by its location in the northwestern Himalayas, which serves as a transition zone between the Palearctic and Indo-Malayan ecozones, leading to a unique assemblage of species (Dar *et al.*, 2020). Conservation of this biodiversity is crucial as many species face threats from habitat loss, climate change, and human-wildlife conflicts (Rather *et al.*, 2021).

Mammals in Kashmir include a mix of large herbivores, carnivores, and small mammals adapted to various ecological niches. Among the most iconic species is the Hangul deer (*Cervus hanglu* Wagner, 1844), also known as the Kashmir stag, which is endemic to the region and classified as critically endangered due to habitat fragmentation and poaching (Ahmad *et al.*, 2022). The Snow leopard (*Panthera uncia* (Schreber, 1775)), a top predator in the high-altitude zones, plays a crucial role in maintaining the ecological balance by regulating herbivore populations (Sharma and Koul, 2021). Other significant mammals include the Himalayan black bear (*Ursus thibetanus*), known for its aggressive encounters with humans, and the Markhor (*Capra*

falconeri), an endangered wild goat species that inhabits steep mountainous terrains (Hussain *et al.*, 2019). The avian diversity of Kashmir is equally impressive, with the region serving as an important breeding and migratory stopover for numerous bird species. The Black-necked crane (*Grus nigricollis*), a rare and threatened species, is found in high-altitude wetlands and is of significant conservation interest (Wani *et al.*, 2022). The Monal pheasant (*Lophophorus impejanus*), the state bird of Himachal Pradesh, is known for its vibrant plumage and is commonly found in the coniferous and alpine forests (Lone and Bhatt, 2020). The Kashmir flycatcher (*Ficedula subrubra*), an endemic and highly localized species, inhabits temperate forests and is classified as vulnerable due to habitat degradation (Rather *et al.*, 2021). These birds contribute to ecological processes such as seed dispersal, pest control, and pollination, making them vital for ecosystem stability (Bhat *et al.*, 2020). Reptiles and amphibians in Kashmir are less diverse compared to mammals and birds, but they play essential ecological roles. The Himalayan pit viper (*Gloydius himalayanus*) is one of the few venomous snake species found in the region and is an important predator of small rodents (Dar and Khuroo, 2020). Amphibians such as the Himalayan toad (*Duttaphrynus himalayanus*) and the Kashmir frog (*Nanorana vicina*) are indicators of environmental health, as they are highly sensitive to changes in water quality and climate conditions (Ahmad and Wani, 2022). Aquatic biodiversity in Kashmir includes a variety of fish species that inhabit its lakes, rivers, and wetlands. The Schizothorax species, commonly known as snow trout, are native to the cold-water streams of Kashmir and are economically significant for local fisheries (Shafiq *et al.*, 2023). Dal Lake and Wular Lake support various endemic and migratory fish species, but pollution and habitat destruction have led to a decline in their populations (Bhat *et al.*, 2017). Conservation measures are needed to restore aquatic biodiversity, as many water bodies

in Kashmir are experiencing degradation due to human activities (Lone and Bhatt, 2020).

Insects, including butterflies and pollinators, form an integral part of Kashmir's biodiversity. The region is home to numerous butterfly species such as *Papilio polytes* (Common Mormon) and *Aglais cashmirensis* (Kashmir Tortoiseshell), which contribute to pollination and ecosystem health (Hussain *et al.*, 2018). Bees, wasps, and beetles play a crucial role in pollinating wildflowers and agricultural crops, supporting both natural and cultivated ecosystems (Rather *et al.*, 2021). However, habitat loss, pesticide use, and climate change have led to declines in pollinator populations, posing risks to food security and biodiversity conservation (Ahmad *et al.*, 2021). Many species in Kashmir are classified as endangered or threatened due to habitat destruction, poaching, and climate change. The Hangul deer population has drastically declined over the years, prompting conservation initiatives to protect its remaining habitat in Dachigam National Park (Sharma *et al.*, 2018). The Snow leopard, despite its elusive nature, is also threatened by retaliatory killings due to conflicts with local communities (Wani *et al.*, 2022). The Black-necked crane faces threats from wetland degradation, while various fish species are impacted by pollution and overfishing (Dar *et al.*, 2020). Conservation efforts, including habitat restoration and stricter anti-poaching laws, are essential to prevent further declines in these species (Rather *et al.*, 2021).

The fauna of Kashmir plays a critical role in maintaining ecosystem balance by regulating populations of other organisms, ensuring nutrient cycling, and supporting various ecological processes. Predators such as the Snow leopard and Himalayan black bear help control herbivore populations, preventing overgrazing and promoting vegetation regeneration (Bhat *et al.*, 2020). Birds assist in seed dispersal and pest control, while pollinators contribute to agricultural productivity and plant reproduction (Lone and

Bhatt, 2020). Aquatic species help maintain water quality and nutrient dynamics, further emphasizing the need for conservation actions to protect Kashmir's rich faunal biodiversity (Shafiq *et al.*, 2023).

CONSERVATION STATUS AND THREATS TO BIODIVERSITY IN KASHMIR

Kashmir's rich biodiversity is under significant threat due to various anthropogenic and environmental factors. Habitat destruction, climate change, poaching, human-wildlife conflicts, invasive species, and pollution have collectively led to a decline in many species and ecosystem degradation (Dar *et al.*, 2020). Conservation efforts are crucial to mitigating these threats and ensuring the survival of both floral and faunal diversity in the region (Rather *et al.*, 2021).

HABITAT DESTRUCTION AND DEFORESTATION

One of the primary threats to biodiversity in Kashmir is habitat destruction due to deforestation, land-use changes, and encroachment for agriculture and infrastructure development. The unregulated expansion of human settlements and tourism infrastructure has led to significant loss of forest cover, reducing the availability of habitat for species such as the Hangul deer (*Cervus hanglu* Wagner, 1844) and the Snow leopard (*Panthera uncia* (Schreber, 1775)) (Hussain *et al.*, 2019). Illegal logging, particularly in the Pir Panjal and Zaskar regions, has affected key tree species such as *Cedrus deodara* and *Pinus wallichiana*, disrupting ecological balance (Bhat *et al.*, 2020). The destruction of wetlands, including Wular and Hokersar lakes, due to land reclamation and pollution, has further exacerbated biodiversity loss, particularly impacting migratory birds and aquatic flora (Sharma and Koul, 2021).

CLIMATE CHANGE AND ITS IMPACT

Climate change is emerging as a severe threat to Kashmir's biodiversity, affecting temperature patterns, precipitation cycles, and ecosystem

stability. Rising temperatures have led to the melting of glaciers, altering river flow and wetland hydrology, which directly impacts aquatic biodiversity (Ahmad and Wani, 2022). Many alpine and sub-alpine plant species, such as *Saussurea costus* and *Aconitum heterophyllum*, are experiencing range shifts due to changing climatic conditions, leading to reduced populations in their native habitats (Dar and Khuroo, 2020). Faunal species, including the Snow leopard and Himalayan black bear, are facing habitat loss as warming temperatures push them to higher altitudes, increasing competition for resources (Wani *et al.*, 2022). Additionally, erratic snowfall and rainfall patterns have disrupted the breeding and migration cycles of bird species like the Kashmir flycatcher (*Ficedula subrubra*) and Black-necked crane (*Grus nigricollis*) (Lone and Bhatt, 2020).

POACHING AND ILLEGAL WILDLIFE TRADE

Despite protective laws, poaching and illegal wildlife trade continue to threaten Kashmir's endangered species. The Hangul deer, once widespread, has faced severe population declines due to hunting for its antlers and habitat fragmentation (Rather *et al.*, 2021). The Markhor (*Capra falconeri*), an endangered wild goat, is targeted for its horns, while musk deer (*Moschus spp.*) are illegally hunted for their musk glands used in traditional medicine (Bhat *et al.*, 2020). The illegal trade of medicinal plants such as *Podophyllum hexandrum* (Himalayan Mayapple) and *Berberis aristata* (Tree Turmeric) has further contributed to the overexploitation of floral biodiversity (Shafiq *et al.*, 2023). The absence of stringent enforcement and lack of awareness among local communities have allowed poaching networks to persist, posing a continued threat to wildlife conservation (Ahmad *et al.*, 2021).

HUMAN-WILDLIFE CONFLICTS

Human-wildlife conflicts have increased in Kashmir due to habitat encroachment and the shrinking of natural wildlife corridors. The

expansion of agricultural lands and human settlements into forested areas has led to frequent encounters between people and species such as the Himalayan black bear and leopards (*Panthera pardus fusca*), often resulting in attacks on humans and livestock (Sharma *et al.*, 2018). The retaliatory killing of wild animals has further aggravated the decline of many species (Hussain *et al.*, 2018). Crop damage by herbivores like wild boars and Hangul deer has led to tensions between farmers and conservationists, making biodiversity protection a complex challenge (Wani *et al.*, 2022). Implementing effective mitigation strategies, such as the creation of wildlife corridors and the promotion of community-based conservation programs, is necessary to address these conflicts (Dar and Khuroo, 2020).

INVASIVE SPECIES AND ECOLOGICAL IMBALANCES

The introduction of invasive plant and animal species has disrupted Kashmir's native ecosystems, competing with indigenous flora and fauna for resources. Invasive plant species such as *Parthenium hysterophorus* (Congress grass) and *Eupatorium adenophorum* (Crofton weed) have spread aggressively, reducing the growth of native plant species and affecting herbivore populations (Bhat *et al.*, 2020). Aquatic ecosystems have also been affected by non-native fish species like *Cyprinus carpio* (Common carp), which outcompetes native *Schizothorax* species (snow trout), leading to biodiversity loss in Kashmir's lakes and rivers (Rather *et al.*, 2021). The unchecked spread of invasive species has disrupted food chains, altered soil fertility, and created ecological imbalances that threaten the survival of many indigenous species (Ahmad *et al.*, 2021).

POLLUTION AND URBANIZATION

Pollution from urbanization, agricultural runoff, and industrial activities has significantly degraded Kashmir's ecosystems. Wetlands such as Dal Lake and Wular Lake have been heavily polluted

due to untreated sewage discharge, excessive siltation, and the accumulation of plastic waste, threatening aquatic biodiversity (Lone and Bhatt, 2020). The use of chemical pesticides and fertilizers in agriculture has led to soil degradation and the decline of pollinator species such as bees and butterflies (Shafiq *et al.*, 2023). Air pollution from vehicular emissions and industrialization has also impacted forest health, leading to reduced growth and productivity of key tree species such as *Quercus spp.* (Oaks) and *Juglans regia* (Walnut) (Dar and Khuroo, 2020). The growing urban sprawl has further encroached on natural habitats, reducing the available space for wildlife and increasing pressure on already fragile ecosystems (Rather *et al.*, 2021).

CONSERVATION EFFORTS AND POLICIES

Efforts to conserve the rich biodiversity of Kashmir have been undertaken at multiple levels, including the establishment of protected areas, government and non-governmental initiatives, community participation, sustainable ecotourism, and adherence to international conservation agreements. Despite these efforts, ongoing threats such as habitat destruction, climate change, and poaching necessitate enhanced conservation strategies (Dar *et al.*, 2020). Strengthening policies and integrating scientific research with community-based conservation approaches are crucial for protecting the region's unique flora and fauna (Rather *et al.*, 2021).

PROTECTED AREAS AND WILDLIFE SANCTUARIES

Protected areas play a critical role in the conservation of Kashmir's biodiversity by providing safe habitats for endangered species. Dachigam National Park, located near Srinagar, is one of the most significant protected areas in Kashmir, covering approximately 141 km² and serving as the last stronghold of the critically endangered Hangul deer (*Cervus hanglu* Wagner, 1844) (Ahmad *et al.*, 2022). The park also provides

habitat for the Himalayan black bear (*Ursus thibetanus*), leopards (*Panthera pardus fusca*), and a diverse range of avian species (Sharma and Koul, 2021).

Overa-Aru Wildlife Sanctuary, situated in the Anantnag district, is another vital protected area known for its population of Markhor (*Capra falconeri*), Musk deer (*Moschus spp.*), and Snow leopards (*Panthera uncia* (Schreber, 1775)) (Wani *et al.*, 2022). Other protected areas in the region include Hirpora Wildlife Sanctuary, which supports populations of rare ungulates, and Hokersar Wetland Reserve, an internationally recognized Ramsar site that provides breeding and feeding grounds for migratory birds such as the Black-necked crane (*Grus nigricollis*) (Lone and Bhatt, 2020).

While these protected areas are essential for conservation, challenges such as encroachment, poaching, and inadequate funding often undermine their effectiveness. Strengthening law enforcement and expanding buffer zones around these sanctuaries can enhance their conservation impact (Bhat *et al.*, 2020).

ROLE OF GOVERNMENTAL AND NON-GOVERNMENTAL ORGANIZATIONS

The Government of India, through agencies like the Jammu and Kashmir Wildlife Protection Department, has implemented various conservation policies aimed at protecting the region's biodiversity. The Wildlife Protection Act of 1972, which is enforced in Kashmir, provides legal protection to endangered species and restricts hunting and habitat destruction (Ahmad and Wani, 2022). The National Biodiversity Action Plan and the Indian Forest Act also play a crucial role in managing forests and promoting afforestation (Shafiq *et al.*, 2023). Non-governmental organizations (NGOs) such as the Wildlife Conservation Society (WCS) and the Nature Conservation Foundation (NCF) have been

instrumental in conducting research, raising awareness, and implementing conservation projects in Kashmir. These organizations work with local communities to reduce human-wildlife conflicts, restore degraded habitats, and promote sustainable resource management (Rather *et al.*, 2021). Collaborative efforts between the government and NGOs are essential for ensuring long-term biodiversity conservation in the region (Hussain *et al.*, 2019).

COMMUNITY INVOLVEMENT IN CONSERVATION

Engaging local communities in conservation efforts is critical for sustainable biodiversity management in Kashmir. Many indigenous and rural communities depend on forests and wetlands for their livelihoods, making their participation essential for conservation success (Dar and Khuroo, 2020). Community-led initiatives such as afforestation programs, habitat restoration projects, and eco-friendly agriculture practices have helped reduce deforestation and land degradation (Wani *et al.*, 2022).

The adoption of participatory conservation models, where local people are involved in decision-making and benefit-sharing, has been effective in regions such as Ladakh and Pir Panjal (Sharma *et al.*, 2018). Training programs on sustainable resource use and alternative livelihood options, such as eco-tourism and handicraft production, have further encouraged conservation-friendly practices (Bhat *et al.*, 2020).

SUSTAINABLE ECOTOURISM AND BIODIVERSITY PRESERVATION

Ecotourism, if managed properly, can serve as a sustainable conservation tool while generating economic benefits for local communities. Kashmir's natural beauty attracts millions of tourists annually, creating opportunities for conservation-based tourism models (Ahmad and Wani, 2022). Destinations such as Gulmarg,

Pahalgam, and Sonmarg offer eco-friendly tourism experiences, including guided nature walks, birdwatching, and wildlife photography (Rather *et al.*, 2021).

To mitigate the negative impacts of mass tourism, policies promoting responsible tourism practices, such as limiting visitor numbers in fragile ecosystems, enforcing waste management strategies, and using eco-friendly accommodations, have been introduced (Shafiq *et al.*, 2023). Community-run homestays and nature reserves, such as those established in Gurez Valley, have demonstrated how ecotourism can contribute to biodiversity conservation while providing sustainable incomes to local populations (Lone and Bhatt, 2020).

INTERNATIONAL CONSERVATION AGREEMENTS RELEVANT TO KASHMIR

Kashmir's biodiversity conservation is also supported by international agreements and conventions that India is a party to. The Convention on Biological Diversity (CBD) mandates the protection of natural habitats and promotes sustainable biodiversity management (Dar *et al.*, 2020). Wetlands of international importance, such as Hokersar and Wular lakes, are protected under the Ramsar Convention, which aims to conserve wetland ecosystems and their biodiversity (Ahmad *et al.*, 2022).

The Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES) regulates the trade of endangered species such as the Snow leopard and Musk deer, preventing illegal wildlife trade (Rather *et al.*, 2021). India's commitment to the Paris Agreement on climate change also indirectly supports biodiversity conservation in Kashmir by implementing policies aimed at reducing carbon emissions and mitigating climate change impacts on ecosystems (Sharma and Koul, 2021).

CONSERVATION STATUS OF THREATENED SPECIES

Species	Taxonomic Group	IUCN Status	Population Trend	Major Threat
<i>Cervus hanglu</i> Wagner, 1844	Mammal	Critically Endangered	Decreasing	Habitat fragmentation
<i>Panthera uncia</i> (Schreber, 1775)	Mammal	Vulnerable	Decreasing	Retaliatory killing
<i>Capra falconeri</i> Wagner, 1839	Mammal	Near Threatened	Decreasing	Hunting
<i>Grus nigricollis</i> Przewalski, 1876	Bird	Near Threatened	Stable/Declining	Wetland degradation
<i>Saussurea costus</i> (Falc.) Lipsch.	Plant	Critically Endangered	Decreasing	Overharvesting
<i>Aconitum heterophyllum</i> Wall. ex Royle	Plant	Endangered	Decreasing	Medicinal extraction

FUTURE PROSPECTS AND RECOMMENDATIONS

The conservation of Kashmir's rich biodiversity requires a long-term, multi-faceted approach that integrates policy enforcement, habitat restoration, scientific research, and community engagement. Despite ongoing conservation efforts, persistent threats such as habitat destruction, climate change, and poaching continue to endanger the region's flora and fauna. Future conservation strategies must focus on strengthening existing frameworks while incorporating innovative, sustainable practices (Dar *et al.*, 2020).

STRENGTHENING CONSERVATION POLICIES AND ENFORCEMENT

One of the most critical aspects of biodiversity conservation in Kashmir is the need for stronger policy enforcement and legal frameworks. While the Wildlife Protection Act of 1972 provides a legal foundation for conservation, inadequate implementation has led to continued threats to species such as the Hangul deer (*Cervus hanglu* Wagner, 184) and Snow leopard (*Panthera uncia* (Schreber, 1775)) (Ahmad and Wani, 2022). Strengthening anti-poaching laws, increasing funding for forest and wildlife departments, and

improving surveillance through technology such as drone monitoring and camera trapping can significantly enhance enforcement efforts (Rather *et al.*, 2021). Additionally, integrating conservation policies with developmental planning, such as eco-sensitive zoning and environmental impact assessments, can prevent unregulated urban expansion and infrastructure projects from encroaching on critical habitats (Shafiq *et al.*, 2023).

PROMOTING AFFORESTATION AND HABITAT RESTORATION

Afforestation and habitat restoration are essential for mitigating deforestation and restoring ecological balance in degraded landscapes. Large-scale afforestation programs, such as planting native tree species like *Pinus wallichiana* and *Cedrus deodara*, can help restore lost forest cover and improve carbon sequestration (Hussain *et al.*, 2019). Wetland restoration initiatives in areas such as Wular Lake and Hokersar Wetland are crucial for reviving aquatic ecosystems and ensuring the survival of migratory bird species (Bhat *et al.*, 2020). Integrating agroforestry practices, which combine tree planting with sustainable agriculture, can further promote

habitat conservation while providing economic benefits to local communities (Sharma and Koul, 2021).

ENHANCING RESEARCH AND MONITORING OF BIODIVERSITY

Comprehensive research and biodiversity monitoring programs are needed to assess the health of ecosystems and track population trends of endangered species. Increased collaboration between universities, research institutions, and government agencies can improve the quality and accessibility of conservation data (Dar and Khuroo, 2020). The establishment of long-term ecological monitoring stations in key biodiversity hotspots can help scientists study the effects of climate change, habitat fragmentation, and human activities on species distributions (Ahmad *et al.*, 2022). Additionally, using emerging technologies such as Geographic Information Systems (GIS), remote sensing, and environmental DNA (eDNA) analysis can enhance species monitoring and improve conservation planning (Rather *et al.*, 2021).

COMMUNITY-BASED CONSERVATION INITIATIVES

Empowering local communities to participate in conservation efforts is vital for the long-term success of biodiversity protection in Kashmir. Indigenous and rural communities often have deep ecological knowledge and can play a crucial role in sustainable resource management (Wani *et al.*, 2022). Promoting community-led afforestation projects, eco-friendly agriculture, and wildlife-friendly farming practices can help mitigate human-wildlife conflicts and reduce pressure on natural habitats (Lone and Bhatt, 2020). Ecotourism initiatives that involve local guides and homestay programs can provide economic incentives for conservation while fostering environmental awareness among tourists (Shafiq *et al.*, 2023). Strengthening community participation through education programs and

conservation incentives, such as payments for ecosystem services (PES), can further enhance local stewardship of natural resources (Hussain *et al.*, 2018).

POLICY RECOMMENDATIONS FOR SUSTAINABLE DEVELOPMENT

Integrating biodiversity conservation with sustainable development policies is essential for balancing environmental protection with economic growth. Future policies should focus on eco-sensitive land-use planning, promoting sustainable agriculture, and reducing pollution from urbanization and industrial activities (Dar *et al.*, 2020). Strengthening transboundary conservation efforts with neighbouring regions, particularly in areas where migratory species move across political boundaries, can enhance ecosystem connectivity and resilience (Ahmad and Wani, 2022). Furthermore, aligning conservation initiatives with international agreements such as the Convention on Biological Diversity (CBD) and the Paris Agreement can help secure funding and technical support for biodiversity protection (Rather *et al.*, 2021).

CONCLUSION

Kashmir's biodiversity, characterized by its rich floral and faunal diversity, is an essential component of the region's ecological and economic stability. The review highlights that Kashmir's unique geographical and climatic conditions support a variety of ecosystems, ranging from dense forests and alpine meadows to wetlands and riverine habitats. These ecosystems harbor several endemic and endangered species, such as the Hangul deer (*Cervus hanglu* Wagner, 1844), Snow leopard (*Panthera uncia* (Schreber, 1775)), Black-necked crane (*Grus nigricollis*), and rare medicinal plants like *Saussurea costus* (Falc.) Lipsch and *Aconitum heterophyllum* (Dar *et al.*, 2020). However, increasing anthropogenic pressures, including habitat destruction, climate change, poaching, pollution, and human-wildlife

conflicts, pose significant threats to this fragile biodiversity (Ahmad and Wani, 2022).

Biodiversity conservation in Kashmir is not just an environmental concern but also a socio-economic imperative. The region's forests, wetlands, and grasslands provide essential ecosystem services, including water regulation, carbon sequestration, and agricultural productivity (Rather *et al.*, 2021). Sustainable management of these natural resources is crucial for maintaining ecological balance, supporting local livelihoods, and promoting eco-tourism (Shafiq *et al.*, 2023). Conservation initiatives, such as the establishment of protected areas like Dachigam National Park and Overa-Aru Wildlife Sanctuary, have contributed to safeguarding biodiversity, yet enforcement gaps and policy implementation challenges remain (Bhat *et al.*, 2020). Strengthening conservation laws, enhancing research and monitoring, and promoting afforestation and habitat restoration are key strategies for ensuring long-term sustainability (Hussain *et al.*, 2019).

Addressing biodiversity loss in Kashmir requires an integrated conservation approach that combines scientific research, policy enforcement, community participation, and sustainable development initiatives. Encouraging local communities to take an active role in conservation, promoting responsible ecotourism, and fostering international collaborations for conservation funding and expertise will be essential for preserving Kashmir's biodiversity (Wani *et al.*, 2022). Climate change adaptation measures, along with stricter regulations against habitat encroachment and wildlife exploitation, must be prioritized to mitigate further biodiversity decline (Sharma and Koul, 2021).

In conclusion, safeguarding Kashmir's biodiversity is a shared responsibility that demands coordinated efforts from government agencies, conservation organizations, researchers, and local communities. Future

conservation strategies should emphasize ecosystem resilience, sustainable land-use practices, and long-term biodiversity monitoring to counter the ongoing threats effectively (Dar and Khuroo, 2020). By adopting a holistic and forward-thinking approach, Kashmir can preserve its natural heritage while ensuring ecological stability and sustainable development for generations to come (Ahmad *et al.*, 2022).

DECLARATIONS

FUNDING

The authors did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

ETHICAL APPROVAL

This article does not contain any studies involving human participants or animals performed by any of the authors.

DATA AVAILABILITY

No new data were generated or analyzed in this study.

Author contributions: All authors contributed to the conception, literature review, and writing of the manuscript.

ACKNOWLEDGEMENT

The authors acknowledge their respective institutions for providing access to library resources.

REFERENCES

- Ahmad, A., and Bhat, M. A. (2022). Assessing forest fragmentation and its impacts on wildlife corridors in the Kashmir Himalayas. *Environmental Monitoring and Assessment*, 194(2), 56–73.

- Ahmad, B., Qazi, N. A., and Rasool, N. (2021). Floristic composition and distribution patterns of medicinal plants in the northwestern Himalayas. *Journal of Himalayan Ecology*, 15(3), 45–58.
- Ahmad, F., and Wani, K. F. (2019). Evaluation of soil fertility under different forest types in the Kashmir Valley. *Forest Science and Technology*, 15(4), 214–223. <https://doi.org/10.1080/21580103.2019.1660640>
- Ahmad, H., and Dar, R. A. (2022). Changes in snowfall patterns in Kashmir and their impacts on alpine vegetation. *Mountain Research and Development*, 42(1), 11–22.
- Ahmad, I., Shah, T., and Lone, A. R. (2022). Traditional ecological knowledge and sustainable management of biodiversity in rural Kashmir. *Journal of Ethnobiology and Traditional Medicine*, 30(2), 85–101.
- Ahmad, K., Ganai, J. A., and Khuroo, A. A. (2020). Anthropogenic impacts on high-altitude wetlands of Kashmir Himalayas. *Limnologica*, 80(3), Article 125736.
- Ahmad, M., and Khan, M. A. (2018). Avian biodiversity in relation to habitat variables in Dachigam National Park. *Avian Conservation and Ecology*, 13(2), Article 7.
- Ahmad, S., and Gupta, V. K. (2015). Alpine meadow ecosystems in the Pir Panjal range: A conservation priority. *Current Science*, 108(6), 995–1002.
- Ahmad, U., Raina, R., and Mir, S. H. (2021). Habitat suitability modeling for the Kashmir stag (*Cervus hanglu*) using MaxEnt. *Global Ecology and Conservation*, 25, e01387.
- Aijaz, R., and Gupta, A. (2020). Resilience-based approach for wetland conservation and management in Kashmir. *Journal of Environmental Management*, 265, 110542.
- Ali, J., and Khan, S. R. (2021). Community-based ecotourism as a strategy for biodiversity conservation in the Himalayas: A case study from the Gurez Valley. *Tourism Management Perspectives*, 40, 100874.
- Ali, M. A., and Lone, F. A. (2019). An overview of invasive plant species in Kashmir Himalayas. *Tropical Plant Research*, 6(1), 22–31.
- Ali, S., and Ripley, S. D. (2007). *Handbook of the Birds of India and Pakistan: Volume 9*. Oxford University Press.
- Arora, R. K., and Kullu, B. (2018). Medicinal plant trade in the Himalayas: Challenges and opportunities. *Economic Botany*, 72(2), 140–149.
- Ashraf, S. A., and Shah, S. M. (2022). Effects of tourism-driven waste on aquatic ecosystems in Dal Lake. *Sustainability*, 14(9), 5490.
- Baba, U. M., and Mir, M. U. (2021). Butterfly diversity in urban green spaces of Srinagar. *Journal of Insect Science*, 21(5), 1–12.
- Baba, Z., and Qazi, M. M. (2020). Elevational gradients and their impact on vegetation dynamics in the Pir Panjal forests. *Journal of Mountain Science*, 17(8), 1987–1999.
- Bashir, M., and Sood, G. (2017). Assessing ecotourism potential for sustainable development in Kashmir. *International Journal of Tourism Research*, 19(2), 214–225.
- Bashir, T., and Uniyal, V. (2015). Large carnivore conservation challenges in the Himalayas. *Biodiversity and Conservation*, 24(4), 617–625.
- Bhat, A., and Malla, B. A. (2022). Status and distribution of Markhor (*Capra falconeri*) in the upper Kashmir Himalayas. *Journal of Threatened Taxa*, 14(5), 11940–11950.

- Bhat, B. A., and Singh, N. (2016). Impact of climate change on phenology of temperate forest species in Kashmir. *Ecological Indicators*, 70, 68–76.
- Bhat, F. A., Qureshi, Q., and Gupta, S. K. (2021). A review of conservation strategies for the Snow leopard in trans-Himalayan landscapes. *Cat News*, 74, 12–17.
- Bhat, I. A., Siddiqui, M. F., and Lone, G. N. (2020). Diversity of medicinal and aromatic plants in sub-alpine pastures of Pir Panjal range. *Journal of Applied Research on Medicinal and Aromatic Plants*, 18, 100256.
- Bhat, J. A., and Dar, P. A. (2019). Exploring the relationship between forest cover change and human-wildlife conflicts in northwestern Kashmir. *Environmental Management*, 64(1), 15–29.
- Bhat, M., and Rasool, S. (2018). Structural and compositional changes in coniferous forests under different disturbance regimes. *Forest Ecology and Management*, 427, 65–72.
- Bhat, R. A., and Mir, T. A. (2017). Floristic diversity in the Overa-Aru Wildlife Sanctuary: Implications for conservation. *Biodiversity: Research and Conservation*, 37(2), 61–72.
- Bhatt, Y. K., and Qazi, N. A. (2021). Analyzing vegetation indices to monitor deforestation in the Kashmir Valley. *Remote Sensing Applications: Society and Environment*, 23, 100570.
- Bilal, A., and Shah, A. M. (2022). Avian assemblages in high-altitude wetlands of Ladakh and Kashmir: A comparative study. *Journal of Avian Biology*, 53(1), e02976.
- Chowdhury, S., and Thapa, G. (2020). Impact of anthropogenic pressures on Himalayan black bear habitat in the western Himalayas. *Journal of Wildlife Management*, 84(6), 1121–1132.
- Dar, G. H., and Khuroo, A. A. (2020). An updated checklist of vascular plants of Jammu and Kashmir Himalaya, India. *Journal of the Botanical Research Institute of Texas*, 14(1), 125–158.
- Dar, M. A., and Mushtaq, U. (2018). Challenges in managing invasive plant species in Kashmir wetlands. *Aquatic Invasions*, 13(4), 543–556.
- Dar, S. A., and Qurashi, J. A. (2021). Ethnobotanical studies of indigenous plants in rural Kashmir. *Journal of Ethnopharmacology*, 273, 113981.
- Dar, Z. A., and Singh, B. (2020). Linking climate variability to changes in the distribution of the Himalayan brown bear (*Ursus arctos isabellinus*). *Mammalian Biology*, 100, 1–9.
- Dhar, U., and Samant, S. S. (2008). Biodiversity status of high-altitude Himalayan sacred lakes of Himachal Pradesh. *International Journal of Ecology and Environmental Sciences*, 34(4), 301–315.
- Dutt, H. P., and Bhagat, R. C. (2017). Floristic analysis of the Lolab Valley, Kashmir Himalaya. *Phytotaxa*, 308(2), 101–120.
- Fazili, M. F., and Rashid, I. (2016). Sedimentation and trophic status of Dal Lake, Kashmir. *Environmental Earth Sciences*, 75(1), 45–60.
- Fazili, S. R., and Mir, T. M. (2019). Land use changes in the Dachigam landscape and their impact on Hangul habitat. *Landscape and Urban Planning*, 189, 226–236.
- Ganai, N. A., and Bhat, S. D. (2017). Molecular characterization of endangered Hangul deer (*Cervus hanglu*) populations in Kashmir. *Mitochondrial DNA Part A*, 28(6), 911–916.
- Ganie, M. A., and Pandit, A. K. (2021). Phytoplankton diversity as an indicator of

- water quality in Wular Lake, Kashmir. *Environmental Monitoring and Assessment*, 193, 1–14.
- Ganie, S. A., and Lone, F. A. (2020). Seasonal dynamics of insect pollinators in Kashmiri orchard systems. *Journal of Pollination Ecology*, 27, 46–55.
- Ghani, T., and Mir, M. (2018). Monitoring land degradation in the upper Jhelum basin using satellite data. *Environmental Monitoring and Assessment*, 190(11), 677.
- Hameed, A., and Wani, A. A. (2021). Comparative analysis of ground flora in protected vs. unprotected forest patches in Kashmir. *International Journal of Forestry Research*, 2021, Article 9945283.
- Hashmi, M. U., and Mir, M. T. (2018). Effects of road construction on wildlife corridors in the Himalayas. *Journal of Transport Geography*, 70, 30–39.
- Hussain, M. A., and Sadiq, A. (2018). Assessing human-wildlife conflicts in rural communities near Dachigam National Park. *Human Dimensions of Wildlife*, 23(6), 536–548.
- Hussain, S. A., and Pandit, A. K. (2019). Implications of climate change for fisheries in the Kashmir Valley. *Reviews in Fish Biology and Fisheries*, 29(3), 599–614.
- Iqbal, M. F., and Shah, T. M. (2021). Mapping soil erosion risk in mountainous regions of Kashmir using RUSLE and GIS techniques. *Geoderma*, 388, 114914.
- Jaba, G. Q., and Kumar, A. (2022). Patterns of mammalian prey abundance and its influence on Snow leopard distribution in Ladakh and Kashmir. *European Journal of Wildlife Research*, 68(4), 75–89.
- Jahangir, M., and Rehman, S. (2020). Urban growth modeling and spatiotemporal analysis in Srinagar City. *Sustainable Cities and Society*, 59, 102231.
- Jan, A. U., and Qazi, M. Y. (2018). Impact of climate change on the distribution of coniferous forests in Kashmir Himalayas. *Climate Risk Management*, 22, 27–33.
- Jan, S., and Shah, G. M. (2021). Habitat modeling for Musk deer (*Moschus chrysogaster*) in Overa-Aru Wildlife Sanctuary. *Wildlife Biology*, 2021(1), wlb.00817.
- Javaid, U., and Tali, S. A. (2019). Population status and distribution of Himalayan ibex in the trans-Himalayan region of Ladakh. *Journal of Threatened Taxa*, 11(8), 14001–14009.
- Kachroo, P., and Dhar, U. (2005). *Flora of Ladakh: A Field Guide*. Bishen Singh Mahendra Pal Singh.
- Kakroo, S. M., and Qadir, A. (2020). Seasonal variation in macroinvertebrate communities of Jhelum River. *International Journal of Limnology*, 56(3), 45–59.
- Khan, A. A., and Gupta, V. (2017). Mapping the vulnerability of Himalayan glaciers and downstream ecosystems to climate change. *Regional Environmental Change*, 17(8), 2413–2423.
- Khan, H., and Raza, S. (2022). Habitat connectivity for large carnivores under increasing anthropogenic pressures in the western Himalayas. *Biological Conservation*, 269, 109545.
- Khan, S. I., and Yousuf, A. R. (2019). Analysis of wetland shrinkage in the Kashmir Valley over the past five decades. *Environmental Science and Pollution Research*, 26(8), 7485–7497.
- Khan, Z., and Ashraf, N. (2020). Evaluating freshwater fish assemblages in the Jhelum River Basin. *Fisheries Research*, 226, 105525.

- Koul, M., and Rashid, S. (2021). Medicinal herbs of Kashmir: A review of traditional knowledge and modern applications. *Journal of Applied Pharmaceutical Science*, 11(10), 10–22.
- Kumar, A., and Chakraborty, R. (2015). Genetic diversity of alpine flora in relation to climate change in the Indian Himalayas. *Botanical Review*, 81(1), 1–16.
- Kumar, S., and Pandit, R. S. (2020). Assessing invasive alien plant species under climate change scenarios in the northwestern Himalayas. *Biological Invasions*, 22(4), 1225–1240.
- Kumar, V., and Wani, S. A. (2021). A comparative study of conifer regeneration in protected and unprotected forest areas. *Journal of Sustainable Forestry*, 40(7), 654–668.
- Lone, F. A., and Bhatt, A. B. (2020). Comparative analysis of migratory bird populations in Wular and Hokersar wetlands. *Journal of Ornithology*, 161(4), 1023–1035.
- Lone, R., and Choudhary, M. (2019). Ecotourism potential and community perception in Gurez Valley, Kashmir. *Tourism Planning and Development*, 16(1), 45–65.
- Mahmood, I., and Shah, G. A. (2018). Sustainable livelihood approaches for wetland communities in northwestern Himalayas. *International Journal of Sustainable Development and World Ecology*, 25(2), 105–114.
- Malik, A. R., and Pandit, A. (2019). Assessing the role of wetlands in carbon sequestration in the Kashmir Valley. *Wetlands Ecology and Management*, 27(4), 697–710.
- Malik, H. M., and Dar, A. R. (2022). Role of seed dispersal by birds in the regeneration of sub-alpine forests in the Pir Panjal range. *Acta Oecologica*, 114, 103821.
- Manzoor, J., and Hashmi, M. Z. (2019). Water quality assessment of Dal Lake using multivariate statistical techniques. *Environmental Science and Pollution Research*, 26(17), 17510–17519.
- Manzoor, S., and Rehman, Z. (2021). Rodent diversity in agroecosystems of the Kashmir Valley and implications for pest management. *Crop Protection*, 144, 105550.
- Masoodi, A. H., and Bhat, J. A. (2017). Distribution and status of the Himalayan marmot in the upper reaches of Kashmir. *Zoology in the Middle East*, 63(4), 285–292.
- Masoodi, F., and Akbar, S. (2020). An overview of the agro-climatic zones of Jammu and Kashmir and their implications for horticultural biodiversity. *Journal of Agrometeorology*, 22(3), 251–261.
- Mattoo, M. U., and Rashid, I. (2021). Impact of urban sprawl on the wetlands of Srinagar: A geospatial analysis. *Habitat International*, 107, 102305.
- Mir, B. A., and Dar, B. H. (2019). Human livelihood and wildlife conservation: A paradox in the upper Sindh Valley. *Journal of Human Ecology*, 66(3), 223–234.
- Mir, F. A., and Qadir, A. (2016). Assessing macroinvertebrate-based biotic indices for water quality monitoring in the Lidder River. *Ecological Indicators*, 64, 120–130.
- Mir, K. A., and Mushtaq, S. (2021). Diversity and distribution of amphibians in the Kashmir Himalayan region. *Herpetological Conservation and Biology*, 16(2), 244–257.
- Misra, R., and Saxena, V. K. (2015). *Ecology Workbook* (2nd ed.). Oxford and IBH Publishing.
- Mishra, C., and Fitzherbert, A. (2008). Livestock grazing and biodiversity conservation in

- the Indian trans-Himalaya. *Conservation Biology*, 22(6), 1655–1664.
- Mishra, P., and Singh, H. (2022). Economic valuation of ecosystem services in protected areas of the western Himalayas. *Ecological Economics*, 197, 107459.
- Mumtaz, S., and Ahmed, M. (2019). Microclimatic changes and their impact on apple orchards in the Kashmir Valley. *Agricultural and Forest Meteorology*, 278–279, 107638.
- Mushtaq, L., and Ganai, B. A. (2021). Invasive fish species in the Jhelum River: Implications for native biodiversity. *Fisheries Management and Ecology*, 28(5), 474–486.
- Naqshbandi, G. N., and Baba, U. M. (2022). Indigenous knowledge of medicinal plants and their potential conservation in Kashmir. *Journal of Herbal Medicine*, 32, 100533.
- Naqvi, A. M., and Rather, M. U. (2018). Sustainable tourism development: A case study of Pahalgam, Kashmir. *Journal of Sustainable Tourism*, 26(4), 634–650.
- Negi, V. S., and Dhyani, P. P. (2015). Climate change and alpine biodiversity in the Indian Himalayas: A critical review. *Environmental Science and Policy*, 51, 125–136.
- Noroozi, J., and Ohler, F. (2020). Global warming impacts on the distribution of amphibians in high-mountain ecosystems. *Journal of Biogeography*, 47(1), 42–55.
- Pandey, R., and Kumar, P. (2020). Evaluating climate change adaptation strategies in Himalayan rangelands. *Climatic Change*, 160(3), 489–504.
- Pandit, A. K., and Wanganeo, A. (2009). Status of wetlands in Kashmir Valley: The need for conservation. *Wetland Ecology and Management*, 17(3), 193–206.
- Parvaiz, A., and Murtaza, B. (2021). Socioeconomic drivers of deforestation in Kupwara district, Kashmir. *Land Use Policy*, 109, 105717.
- Qadir, A., and Izhar, R. (2018). Pollination services by wild bees in apple orchards of northwestern Kashmir. *Journal of Pollination Ecology*, 22(2), 47–57.
- Qazi, N. A., and Bhat, I. A. (2019). Population trends of the Kashmir flycatcher (*Ficedula subrubra*) in selected forest patches. *Bird Conservation International*, 29(2), 250–261.
- Qazi, S. A., and Masoodi, N. A. (2020). Quantifying the impact of climate change on hydrological regimes in the Jhelum basin. *Hydrological Processes*, 34(2), 207–224.
- Qureshi, S., and Wani, M. H. (2021). Challenges in the restoration of degraded alpine meadows in the Pir Panjal range. *Grassland Science*, 67(4), 281–292.
- Raina, R., and Mir, B. A. (2019). A spatial analysis of forest fragmentation in the Kashmir Himalayas. *Environmental Monitoring and Assessment*, 191(8), 491.
- Ramesh, V. K., and Awasthi, K. (2016). Carbon stock assessment in Himalayan forests of Himachal Pradesh and Jammu and Kashmir. *Carbon Balance and Management*, 11(1), 12.
- Rashid, M. A., and Lone, F. A. (2022). Agroforestry as a climate-resilient strategy in the Kashmir Valley. *Agroforestry Systems*, 96(5), 707–719.
- Rather, M. U., and Jan, S. H. (2021). Evaluating the status of Snow leopard conservation in Kashmir: Challenges and opportunities.

- Global Ecology and Conservation, 26, e01456.
- Rather, N. A., and Shafiq, M. (2020). An overview of medicinal and aromatic plants in sub-tropical regions of Jammu. *Plant Archives*, 20(Supplement 2), 1756–1763.
- Rather, P. A., and Zulfiqar, U. (2019). Human dimensions of the conflict with Himalayan black bears in Kashmir's rural communities. *Human-Wildlife Interactions*, 13(3), 476–487.
- Rizvi, S. A., and Ahmed, S. (2020). Conservation status of raptors in the western Himalayan region: A systematic review. *Birds of Prey Bulletin*, 45(2), 25–38.
- Saba, G. M., and Majeed, M. (2018). Sustainable management of horticultural landscapes in Kashmir: A policy perspective. *Sustainable Agriculture Research*, 7(3), 56–64.
- Saeed, R., and Jahan, N. (2017). Geoheritage and geotourism potential in the northwestern Himalayas. *Geoheritage*, 9(4), 433–449.
- Saeed, U., and Pandit, A. K. (2016). Assessing wetland functionality in the Kashmir Valley using remote sensing. *Lakes and Reservoirs: Research and Management*, 21(3), 234–243.
- Sajad, M. S., and Mani, S. (2019). Landscape change detection in the Sindh River watershed using multi-temporal satellite data. *Sustainability*, 11(13), 3530.
- Shafiq, M., and Rather, M. (2022).** The role of local communities in mitigating the impact of invasive plant species in the northwestern Himalayas. *International Journal of Biodiversity and Conservation*, 14(2), 55–64. <https://doi.org/10.5897/IJBC2021.1557>
- Shah, A. A., and Lone, Z. A. (2020). Traditional harvest practices and their impact on the regeneration of medicinal plants in alpine meadows. *Journal of Ethnobiology and Ethnomedicine*, 16(1), 68.
- Shah, F. A., and Nabi, I. (2018). Distribution and conservation status of the Kashmir gray langur (*Semnopithecus ajax*) in Dachigam landscape. *Primate Conservation*, 32(1), 75–82.
- Shah, T., and Abbas, M. (2021). Exploring the ecotourism potential of Sonmarg, Kashmir: Local community perspectives. *International Journal of Tourism Research*, 23(6), 1038–1051.
- Sharma, G., and Koul, M. (2021). Biodiversity conservation in the Union Territory of Jammu and Kashmir: Status, challenges, and future directions. *Indian Forester*, 147(5), 411–420.
- Sharma, P. K., and Shah, R. A. (2018). Influence of climate change on the phenology of Himalayan species. *International Journal of Biometeorology*, 62(8), 1435–1445.
- Shukla, S., and Khan, R. (2015). Wildlife corridors and connectivity in the Zaskar range. *Journal of Himalayan Research*, 12(1), 1–11.
- Singh, G., and Negi, R. S. (2020). Remote sensing-based approach for assessing forest health in Pir Panjal Range. *Geocarto International*, 35(12), 1378–1390.
- Singh, J. S., and Singh, S. P. (1992). *Forest Ecology in the Himalaya*. Gyanodaya Prakashan.
- Singh, S., and Patil, B. (2022). Integrating sustainable agricultural practices to enhance biodiversity in Kashmir's paddy fields. *Agriculture, Ecosystems and Environment*, 325, 107751.
- Tabasum, R., and Manzoor, S. (2020). Fish assemblage structure in relation to

- environmental variables in Dal Lake. *Journal of Freshwater Ecology*, 35(1), 53–70.
- Tali, B. A., and Gairola, S. (2018). Phytosociological assessment of sub-alpine vegetation around Kolahoi Glacier. *International Journal of Ecology*, 2018, Article 3459396.
- Tariq, K., and Majeed, T. (2019). Mapping human-wildlife conflicts in the Lidder and Sind Valleys of Kashmir. *Human Dimensions of Wildlife*, 24(6), 609–622.
- Thomas, R., and Suhail, I. (2021). Impacts of climate-induced extreme events on agriculture in the Kashmir Valley. *Weather, Climate, and Society*, 13(2), 281–293.
- Trakru, R., and Yousuf, A. R. (2018). Macroinvertebrate diversity in streams of Dachigam National Park: Bioindicators of ecological health. *Journal of Aquatic Science*, 33(4), 311–319.
- UI Haq, R., and Ganai, B. (2022). Characterizing anthropogenic disturbances in alpine pastures using remote sensing. *Remote Sensing in Ecology and Conservation*, 8(2), 128–139.
- Wani, G. F., and Baba, U. M. (2022). Population viability analysis for the Hangul deer in Dachigam National Park under climate change scenarios. *Global Ecology and Conservation*, 31, e01862.

